

Dynamics of Perturbed Relative Equilibria of Point Vortices on the Sphere or Plane

G. W. Patrick

Department of Mathematics and Statistics
University of Saskatchewan
Saskatoon, Saskatchewan, S7N 5E6
Canada

June 1999

Abstract

Stable assemblages of localized vortices exist which have particle-like properties, such as mass, and which can interact with one another when they closely approach. In this article I calculate the mass of these localized states and numerically investigate some aspects of their interactions.

Introduction

The system of N point vortices on a sphere is a Hamiltonian system with symmetry $SO(3)$. Steadily rotating arrangements of vortices are the relative equilibria of these systems, and the nonlinear stability of these relative equilibria may be analyzed as a matter of routine, at least in principle. If stable then such an arrangement of vortices will, under sufficiently small perturbation, approximately maintain its “shape”, so that its subsequent motion will be nearly its initial condition up to the action of the symmetry group $SO(3)$. However, if the momentum of the relative equilibrium (i.e. its center of vorticity) is zero then the position of such a relative equilibrium on the sphere is not stable under perturbation to nonzero momentum, as has been established in Patrick [1992, 1995, 1998].

In this article I show that there are arbitrarily localized relative equilibria of N vortices with zero momentum, and that these relative equilibria are formally stable for $N = 3$ and $N = 4$. Under small perturbation to nonzero momentum, they move on the sphere, to first order in the momentum perturbation, as a free particles move under the influence of a magnetic monopole. The effect of the monopole diminishes as the relative equilibria become more localized. Since a given momentum perturbation implies a particular velocity of motion of each relative equilibrium as a whole, they have mass, and using a perturbation theory of Patrick [1995], I find a formula for that mass.

For $N = 4$, when two such relative equilibria are placed on the sphere, they will move independently until they approach within a distance comparable to their size, whereupon an interaction occurs. For example, one might collide with its opposite (obtained by reversing the first's component vortex strengths), the result being "radiation" in the form of dipole pairs, and generally opposites repel and likes attract. Here in this article I numerically determine some general features of these interactions, which are the subject of ongoing investigations.

Over small regions of the sphere, the point vortex system on the sphere becomes the point vortex system on the plane, and since the above relative equilibria may be arbitrarily localized, one expects such in the planar case as well. Towards the end of this article I show this in indeed the case. The planar system of N point vortices will be a worthy motivating example for extensions of the stability theory of Hamiltonian systems with symmetry to the case of non-equivariant momentum mappings and/or noncompact groups.

1 Context and Notations

I briefly summarize here the basic elements of the system of N point vortices on the sphere and on the plane. For more details, see Kidambi and Newton [1998], Pekarsky and Marsden [1998], and the many references therein.

The system of N point vortices of strengths $\Gamma_i \in \mathbb{R}$ on the sphere S^2 of radius R is the Hamiltonian system with phase space $P^{\text{sp}} \equiv (S^2)^N$ and Hamiltonian

$$H^{\text{sp}}(x_1, \dots, x_N) \equiv \frac{1}{4\pi R^2} \sum_{m < n} \Gamma_n \Gamma_m \ln l_{mn}^{-2}, \quad (1)$$

where x_n represents the location of the n^{th} vortex and $l_{mn}^2 = 2(R^2 - x_m \cdot x_n)$ is the chord length between x_n and x_m . The symplectic structure ω^{sp} on P^{sp} is a direct sum of the symplectic structures on S^2 weighted by vortex strengths; specifically

$$\omega^{\text{sp}} \equiv \bigoplus_{n=1}^N \frac{\Gamma_n}{R} \omega_{S^2}, \quad (2)$$

where

$$\omega_{S^2}(x)(v, w) \equiv -\frac{1}{R^2} x \cdot (v \times w). \quad (3)$$

Most significant from a geometric mechanics point of view is that this system admits the symmetry of diagonal multiplication of $SO(3)$ on P^{sp} with momentum map

$$J^{\text{sp}} \equiv -\frac{1}{R} \sum_{n=1}^N \Gamma_n x_n. \quad (4)$$

These notations are as in Pekarsky and Marsden [1998].

In the system of N vortices in the plane the n^{th} vortex has location $z_n = x_n + iy_n \in P^{\text{pl}} \equiv (\mathbb{C}^2)^N$, and the Hamiltonian and symplectic form are

$$H^{\text{pl}} \equiv -\frac{1}{4\pi} \sum_{m < n} \Gamma_n \Gamma_m \ln |z_n - z_m|^2, \quad \omega^{\text{pl}} \equiv \bigoplus_{n=1}^N \Gamma_n \omega_0, \quad (5)$$

where generally I will use the notation $\omega_0(a, b) \equiv -\text{Im}(a\bar{b})$ for complex numbers $a, b \in \mathbb{C}$. This system admits the symmetry group $SE(2) = \{(e^{i\theta}, a)\}$ of Euclidean symmetries acting diagonally on each factor $\mathbb{C}$ of P^{pl} by $(e^{i\theta}, a) \cdot z \equiv e^{i\theta}z + a$, and a momentum mapping is

$$J^{\text{pl}} \equiv -\sum_{n=1}^N \Gamma_n \begin{bmatrix} \frac{1}{2}|z_n|^2 \\ iz_n \end{bmatrix}, \quad (6)$$

where $\mathfrak{se}(2)^*$ is identified with $\mathfrak{se}(2) = \{(\dot{\theta}, \dot{a})\} \cong \mathbb{R}^3$ by the standard inner product of $\mathbb{R}^3$. These notations are as in Lewis and Ratiu [1996].

When attention is confined to a sufficiently small region of the sphere, the spherical system becomes the planar one. For example, using on each factor the map

$$\phi(z) = (z, \sqrt{R^2 - |z|^2}) \quad (7)$$

to pull back the sphere to the plane one obtains

$$\phi^* \omega^{\text{sp}} = -\frac{1}{zR} \omega^{\text{pl}} \approx -\frac{1}{R^2} \omega^{\text{pl}}, \quad \phi^* H^{\text{sp}} \approx -\frac{1}{R^2} H^{\text{pl}}. \quad (8)$$

Thus the pulled back equations of motion are approximately the same as the equations of motion of the planar system, since the factors of $-1/R^2$ here multiply both the symplectic form and the Hamiltonian.

As noted in Adams and Ratiu [1988], the momentum mapping of the planar system is equivariant if and only if the total vortex strength $\sum_n \Gamma_n$ vanishes. Indeed, the adjoint and coadjoint actions of $SE(2)$ are

$$\begin{aligned} \text{Ad}_{(e^{i\theta}, a)}(\dot{\theta}, \dot{a}) &= (\dot{\theta}, e^{i\theta} \dot{a} - i\dot{\theta}a), \\ \text{CoAd}_{(e^{i\theta}, a)}(\mu, \nu) &= (\mu - \omega_0(e^{i\theta}\nu, a), e^{i\theta}\nu), \end{aligned} \quad (9)$$

and the deviation of the momentum map J^{pl} from equivariance is the cocycle $\sigma : SE(2) \rightarrow \mathfrak{se}(2)^*$ defined by $\sigma \equiv J^{\text{pl}}((e^{i\theta}, a) \cdot p) - \text{Coad}_{(e^{i\theta}, a)} J^{\text{pl}}(p)$ (the evaluation point $p \in P$ is irrelevant). For the case at hand the cocycle is

$$\sigma(e^{i\theta}, a) = -(\sum_n \Gamma_n) \begin{bmatrix} \frac{1}{2}|a|^2 \\ ia \end{bmatrix},$$

and the derivative of this cocycle at the identity is the skew-symmetric two form $\Sigma : \mathfrak{se}(2)^2 \rightarrow \mathbb{R}$ given by

$$\Sigma((\dot{\theta}_1, \dot{a}_1), (\dot{\theta}_2, \dot{a}_2)) = (\sum_n \Gamma_n) \omega_0(\dot{a}_1, \dot{a}_2). \quad (10)$$

Generalities on non-equivariant momentum mappings may be found in Abraham and Marsden [1978]; a prime fact is the momentum commutation identity

$$\{J_\xi, J_\eta\} = J_{[\xi, \eta]} - \Sigma(\xi, \eta). \quad (11)$$

Both the planar and spherical systems have simple closed form solutions in the case $N = 2$: for the sphere any two vortices evolve as the action of the one parameter group with generator $J^{\text{sp}}/2\pi l_{12}^2$ while for the plane they evolve as the action of the one parameter group with generator $\dot{\theta} = (\Gamma_1 + \Gamma_2)/2\pi|z_1 - z_2|^2$, $\dot{a} = -J^{\text{pl}}/2\pi|z_1 - z_2|^2$. Numerically the action of these one parameter subgroups is easily computed, and since the Hamiltonians H^{sp} and H^{pl} are sums of pairwise interactions, the full system of N vortices may be numerically integrated in a symplectic, symmetry preserving and momentum preserving way using splitting methods, as in Channell and Neri [1993].

2 The Relative Equilibria

To find the relative equilibria for the system of N vortices in the sphere one seeks $\xi_e \in \mathfrak{so}(3) \cong \mathbb{R}^3$ and $p_e \in P^{\text{sp}}$ such that $dH^{\text{sp}}(p_e) - dJ_{\xi_e}(p_e) = 0$. Equivalently, using the obvious extensions of H^{sp} and J^{sp} to $(\mathbb{R}^3)^N$, one can solve the equations

$$\nabla_m H^{\text{sp}} - \nabla_m J_{\xi_e}^{\text{sp}} = \tilde{\lambda} \nabla \left(\frac{1}{2} |x_m|^2 \right) = \tilde{\lambda}_m x_m, \quad m = 1, \dots, N,$$

where the $\tilde{\lambda}_m \in \mathbb{R}$ are Lagrange multipliers. Inserting (1) and (4) yields the N equations

$$\sum_{n \neq m} \frac{\Gamma_n}{l_{nm}^2} = \lambda_m x_m + 2\pi R \xi_e, \quad \lambda_m \equiv -\frac{2\pi R^2}{\Gamma_m} \tilde{\lambda}_m, \quad m = 1, \dots, N. \quad (12)$$

General analytic solutions are not to be expected to these nontrivial nonlinear equations.

I seek a manifold of solutions to (12) contained in the zero level set of J^{sp} , which in some limit is confined to arbitrarily small regions of phase space, and which is formally stable near that limit. I try a regular polygonal configuration where the first $N - 1$ vortices of equal strengths surround the (possibly different strength) N^{th} central vortex, $N \geq 3$. I will denote the strength of the central vortex by Γ . For convenience I locate the central N^{th} vortex at $R\mathbf{k}$ where $\mathbf{k} \equiv (0, 0, 1)$, and the first vortex at $(R \sin \alpha, 0, R \cos \alpha)$, so that α is the opening angle—the angle at the center of the sphere between any outer vortex and the central vortex. The momentum is zero if and only if

$$\Gamma_1(x_1 + x_2 + \dots + x_{N-1}) + \Gamma x_N = (N - 1)\Gamma_1 R \cos \alpha \mathbf{k} + \Gamma R \mathbf{k} = 0$$

which is equivalent to

$$\Gamma_1 = -\frac{\Gamma}{(N - 1) \cos \alpha}, \quad (13)$$

while equations (12) reduce to $\lambda_1 = \dots = \lambda_{N-1}$ and the two equations

$$\Gamma_1 X_0 + \frac{R\Gamma}{l_{1N}^2} \mathbf{k} = \lambda_1 x_1 + 2\pi R \xi_e, \quad (14)$$

$$\frac{(N-1)\Gamma_1 R \cos \alpha}{l_{1N}^2} \mathbf{k} = \lambda_N R \mathbf{k} + 2\pi R \xi_e, \quad (15)$$

where ξ_e depends on Γ , N , α and R , and where

$$X_0 \equiv \frac{x_2}{l_{21}^2} + \frac{x_3}{l_{31}^2} + \dots + \frac{x_{N-1}}{l_{N-1,1}^2}. \quad (16)$$

Setting

$$\mu_N \equiv \sum_{k=1}^{N-1} \frac{1}{e^{-2\pi i k/N} - 1}, \quad \nu_N \equiv \sum_{k=1}^{N-1} \frac{1}{|e^{2\pi i k/N} - 1|^2},$$

so that

$$\mu_N + \nu_N = \sum_{k=1}^{N-1} \frac{e^{2\pi i k/N}}{|e^{2\pi i k/N} - 1|^2},$$

one sees by symmetry that μ_N and ν_N are real, and also that

$$\mu_N = -\frac{1}{2}(N-1), \quad \nu_N = \frac{1}{12}(N^2-1).$$

Then scaling μ_{N-1} and ν_{N-1} gives

$$X_0 \cdot \mathbf{i} = \frac{\mu_{N-1} + \nu_{N-1}}{R \sin \alpha}, \quad X_0 \cdot \mathbf{j} = 0, \quad X_0 \cdot \mathbf{k} = \frac{\nu_{N-1} \cos \alpha}{R \sin^2 \alpha}. \quad (17)$$

Equation (15) implies ξ_e is along $\mathbf{k}$ and in (14) it is clear that the $\mathbf{j}$ component of both sides is zero, so writing (14) and (15) in components gives three linear equations in the three unknowns λ_1 , λ_N , and $\xi_e \cdot \mathbf{k}$. These equations can be routinely solved to obtain

$$\lambda_1 = -\frac{\Gamma}{R^2 \cos \alpha \sin^2 \alpha} \frac{(N-2)(N-6)}{12(N-1)}, \quad (18)$$

$$\lambda_N = -\frac{\Gamma}{R^2 \sin^2 \alpha} \left(\cos \alpha + \frac{N}{2(N-1)} \right), \quad (19)$$

$$\xi_e \cdot \mathbf{k} = \frac{\Gamma}{4\pi R^2 \sin^2 \alpha} \left(\frac{1}{N-1} + \cos \alpha \right). \quad (20)$$

In the limit $\alpha \rightarrow 0$, from (13), the total vortex strength

$$\sum_n \Gamma_N = \Gamma + (N-1)\Gamma_1 = \Gamma \left(1 - \frac{1}{\cos \alpha} \right) \quad (21)$$

vanishes while the regular polygon of outer vortices collapses upon the central vortex. Although these relative equilibria depend on α , Γ and N , I will denote them below simply by p_e .

Formal stability means definiteness of the Hessian of the reduced Hamiltonian at the relative equilibrium, and, by the energy-momentum, method as in Marsden and Ratiu [1994], this is equivalent to definiteness of the function $H^{\text{sp}} - J_{\xi_e}$ on a subspace tangent to the momentum level set and complimentary to the tangent space to the group orbit. For $N = 3$ the reduced spaces are points and formal stability is immediate. I have determined the stability p_e for $N = 4, 5, 6$ with the aid of a symbolic manipulator. For $N = 4$ the relative equilibria are formally stable without conditions, and in particular formally stable arbitrarily near the limit $\alpha \rightarrow 0$. For $N = 5$, but numerically now, one has formal stability if and only if $\alpha > 1.951 \approx 111.8^\circ$ and for $N = 6$ if and only if $\alpha > 2.245 \approx 128.6^\circ$. Presumably this pattern continues and the relative equilibria are formally stable for $N > 6$ if and only if α is sufficiently near π .

As already mentioned, since the momenta of such relative equilibria are zero, by Patrick [1992], formal stability implies stability only modulo $SO(3)$; dynamically, these relative equilibria, when perturbed to nonzero momentum, approximately maintain their shape, which oscillates on a fast time scale, while they move around the sphere on a slow time scale. The left of Figure (1) is the result of a simulation and shows a typical motion of one such relative equilibrium. For short, I will call a point of phase space that results from perturbing one of the relative equilibria p_e a “preq”.

The motion of preq may be approximated as a direct application of the perturbation theory of Patrick [1995]. In summary, the symplectic $SO(3)$ symmetry implies that the linearization of the relative equilibrium has double eigenvalues and hence can be expected to have a nonzero nilpotent part, say N_α . It can be shown that the image of N_α is contained in the tangent space to the group orbit and the tangent space to the momentum level set is contained in the kernel of N_α . Consequently there is a unique (it can be shown to be symmetric) bilinear form (here with the same name) N_α on $\mathfrak{so}(3)^*$ such that the following diagram commutes

$$\begin{array}{ccc} T_{p_e} P^{\text{sp}} & \xrightarrow{N_\alpha} & T_{p_e} P^{\text{sp}} \\ \downarrow dJ(p_e) & & \uparrow \xi \mapsto \xi p_e \\ \mathfrak{so}(3)^* & \xrightarrow{N_\alpha} & \mathfrak{so}(3) \end{array}$$

I have calculated the N_α in the case $N = 3$ and $N = 4$; they are the diagonal matrices with entries J_{N1}, J_{N2}, J_{N3} where

$$J_{31} = J_{32} \equiv \frac{1}{8\pi R^2 \cos \alpha}, \quad (22)$$

$$J_{33} \equiv \frac{1}{8\pi R^2 \sin^4 \alpha} (2 \cos^3 \alpha + 3 \cos^2 \alpha + 2 \cos \alpha - 1), \quad (23)$$

and

$$J_{41} = J_{42} \equiv -\frac{1}{2\pi R^2} \frac{3 \cos^2 \alpha + \cos \alpha + 2}{(1 - \cos \alpha)(9 \cos^2 \alpha + 4 \cos \alpha + 3)}, \quad (24)$$

$$J_{43} \equiv \frac{1}{12\pi R^2 \sin^4 \alpha} (3 \cos^3 \alpha + 4 \cos^2 \alpha + 3 \cos \alpha - 2). \quad (25)$$

The perturbation theory asserts that to first order the motion of the preq is that of the *drift system*: the reduction, by the *normal form symmetry* of the right-hand action of the torus generated by $\mathbf{k} \in \mathfrak{so}(3)$, of the left invariant *drift Hamiltonian* $\pi \mapsto \pi \cdot \xi_e + \frac{1}{2} \pi^t N_\alpha \pi$ on $T^*SO(3)$ with canonical symplectic form. That the drift Hamiltonian is invariant under this toral action is predicted by the theory and is evident by the equality $J_{N1} = J_{N2}$. This flavor of the theory is subject to the caveat the the frequencies of the linearization of the equilibrium are not in 1-1 resonance with the rotation frequencies of the relative equilibrium itself; were there to be such a 1-1 resonance the preq would acquire the structure of a magnetic moment, as in Patrick [1998]. For the case $N = 3$ this is not an issue since the reduced spaces are points and so there are no reduced frequencies. For $N = 4$ one verifies by calculating the linearization that if the reduced frequencies are $\pm \omega_{\text{red}}$ then

$$\frac{|\xi_e|^2}{\omega_{\text{red}}^2} = 1 - 3 \frac{(9 \cos^2 \alpha + 4 \cos \alpha + 3) \sin^2 \alpha}{3 \cos^2 \alpha + 2 \cos \alpha + 3}.$$

Since the second term does not vanish for $0 < \alpha < \pi$, the group frequencies $\pm \xi_e$ are never equal to the reduced frequencies and the simpler perturbation theory suffices.

If the location of the preq on the sphere is denoted by y , then the equations of motion for it (i.e. the equations of motion of the drift system) turn out to be

$$m_\alpha \frac{d^2 y}{dt^2} = -m_\alpha \frac{|v|^2}{R} \frac{y}{R} + \sigma B \times v, \quad v \equiv \frac{dy}{dt}, \quad B \equiv \frac{1}{R^3} y, \quad (26)$$

where σ the momentum associated to the normal form symmetry and

$$m_\alpha \equiv -\frac{1}{J_{N1} R^2}.$$

Thus, the drift system is the same as that of particle of mass m_α and charge σ moving on the sphere under the influence of the magnetic monopole B . Given that one is perturbing a relative equilibrium where the central vortex has been located at $R\mathbf{k}$, the initial location of the preq may be taken to be $y(0) = R\mathbf{k}$, while its initial velocity $v(0)$ and the charge σ may be obtained from the momentum perturbation $\Delta\mu$ by the equations

$$v(0) = \frac{1}{m_\alpha R^2} y(0) \times \Delta\mu, \quad \sigma = -\Delta\mu \cdot \mathbf{k}.$$

By solving Equations (26), one finds that the prediction of the theory is that the preq will rotate about the perturbed momentum, say $\Delta\mu$, with angular velocity $J_{N1} \Delta\mu$.

Specified momentum perturbations of the relative equilibria p_e may be accomplished by moving the central vortex from its original location at $R\mathbf{k}$ to $(R \sin \delta, 0, R \cos \delta)$ while changing the angle α to $\alpha + \Delta\alpha$, as follows. The momentum of the perturbed configuration is easily verified to be

$$J^{\text{sp}} = -(n-1)\Gamma_1 \begin{bmatrix} 0 \\ 0 \\ \cos(\alpha + \Delta\alpha) \end{bmatrix} - \Gamma \begin{bmatrix} \sin \delta \\ 0 \\ \cos \delta - 1 \end{bmatrix},$$

so this is $\Delta\mu \equiv [\Delta\mu_1, 0, \Delta\mu_3]$ if

$$\sin \delta = -\frac{\Delta\mu_1}{\Gamma}, \quad (27)$$

$$\cos(\alpha + \Delta\alpha) - \cos \alpha = -\frac{1}{(N-1)\Gamma_1}(\Delta\mu_3 + \Gamma(\cos \delta - 1)), \quad (28)$$

and arbitrary directions in $\Delta\mu$ may be obtained by rotating this. As these formulas are to be used to perturb a relative equilibrium it is understood that $\Delta\mu_1$ and $\Delta\mu_3$ are small; just how small depends on the validity of the approximation that is the drift system. The momentum of the relative equilibrium, being zero, does not itself provide a scale. Certainly, though, the perturbation should not significantly affect the geometry of the relative equilibrium, meaning it should not displace the vortices by amounts comparable to the diameter 2α . In the using of (27) and (28) one should therefore ensure

$$\left| \frac{\delta}{\alpha} \right| \ll 1, \quad \left| \frac{\Delta\alpha}{\alpha} \right| \ll 1 \quad (29)$$

Using (27) and (28) I have simulated the 4-vortex preq corresponding to $\alpha = \pi/6$ on the unit sphere for momenta $\Delta\mu$ small multiples of the vector $[2, 0, 3]$, and calculated the angular velocities of the preq. The results are on the right of Figure (1). The drift rates fit well to a cubic with slope $|J_{41}|$ at zero.

An essential aspect of the preq is that their location is ill defined as a concept. Suppose for example one assigns the location $R\mathbf{k}$ to a preq corresponding to the state $p_0 \in P^{\text{sp}}$ at some particular time, and sometime later the state of the system is $p_1 \in P^{\text{sp}}$. If there is a group element $A \in SO(3)$ such that $Ap_0 = p_1$ then the location corresponding to p_1 is $RA\mathbf{k}$, unequivocally. However, for $N \geq 4$ the reduced phase space for the vortex system has dimension $2N - 4 \geq 4$, so the flow on this reduced phase space is usually at least as complicated as a toral flow with two incommensurate frequencies. Thus the reduced flow may never repeat itself, and there may never be an A such that $Ap_0 = p_1$ exactly. For the simulations above the location of the preq was decreed to be the average of the locations of its constituent vortices; another possibility for example is the location of its central vortex. The preq alters its shape on a fast time scale compared with its overall motion, and this gives a statistical character to the meaning of its “location” on the sphere. The location of the preq becomes ever more exact and its character ever less statistical as the perturbation falls to zero.




Figure 1: Left: A 4-vortex preq with opening angle $\alpha = \pi/6$ subjected to a large perturbation subsequently moves on the sphere in a circular path about the perturbed momentum, which points towards you. The sense of rotation as is clockwise, the reverse of the right hand sense obtained from the perturbed angular momentum. Shown are the paths of the central vortex and one outer vortex; the paths of the other two outer vortices are suppressed to avoid cluttering the picture. Upper right: the drift rate vs $|\Delta\mu|$; the curve shown is a cubic with slope at 0 equal to J_{41} . Lower right: the angular velocity of a preq divided by $|\Delta\mu|$, calculated over 200 consecutive intervals of its path.

This problem is illustrated in the bottom right of Figure (1), where path of a preq has been regularly sampled and plotted are the values of J_{41} from angular velocities obtained from consecutive changes in the angle that the preq makes with its initial condition. Were the preq to exactly follow the drift system the plot would be constant, but instead one gets a variation about the average value of 0.4602. This average is just .07% off the value of $J_{41} = .4599$ predicted by Equation (24).

The issue of whether it is possible to define the location of a preq may be put more deeply as follows. Let the Marsden-Weinstein reduced space for the vortex system at its zero momentum level be P_0^{sp} with reduced Hamiltonian $\bar{H}^{\text{sp}}$; the relative equilibrium p_e corresponds to the equilibrium, say $\bar{p}_e$ of the reduced system. In Patrick [1995] it is shown that there is a symplectomorphism defined near the group orbit $SO(3) \cdot p_e$ and onto a neighborhood of $\bar{p}_e$ times the zero section of $T^*SO(3)$ such that the Hamiltonian H^{sp} becomes

$$\tilde{H}^{\text{sp}}(x, \pi) \equiv \bar{H}^{\text{sp}}(x) + \pi \cdot \xi_e + \frac{1}{2} N_\alpha \pi^2 + \text{h.o.t.}$$

Thus the reduced degrees of freedom are linked to the overall motion by the higher order terms depending on both x and π , and if these terms vanish then the location of the preq may be unequivocally assigned by following the fiber of $T^*SO(3)$ to its zero section. Conversely, the location of the preq is ill defined in as much as one is obstructed in removing these higher order terms.

3 Small Opening Angles

With opening angles as large as $\alpha = \pi/6$, such as in Figure (1), the relative equilibrium p_e does not have the appearance of a localized particle. But one may choose α arbitrarily small, and now I discuss some aspects of small α and report some numerics for $\alpha = \pi/2048 \approx .088$ deg.

Firstly, in the $\alpha \rightarrow 0$ limit there is an essential difference between the 3-vortex and 4-vortex preq: for the 3-vortex preq, from (22),

$$m_\alpha = -8\pi + O(\alpha^2),$$

while for the 4-vortex preq, from (24),

$$m_\alpha = \frac{8}{3}\pi\alpha^2 + O(\alpha^4).$$

Since the mass of the 3-vortex preq does not fall with α while that of the 4-vortex preq does, the 3-vortex preq is very heavy in comparison to the 4-vortex preq for small α . Thus, large momentum perturbations are required to move a 3-vortex preq. As shall be shown immediately, momentum perturbations must also fall with α , or else destroy the relative equilibrium by too grossly perturbing it. Thus, for dynamical purposes the 3-vortex preq is infinitely heavy in the $\alpha = 0$ limit. This is confirmed upon passage to the planar system, as will be seen in Section (4).

The restrictions (29) in the $\alpha = 0$ limit have a different character as they affect $\Delta\mu_1$ vs. $\Delta\mu_3$. The effect on $\Delta\mu_1$ is clear: directly from (27) and the first of (29)

$$|\Delta\mu_1| = |\Gamma \sin \delta| \ll O(\alpha).$$

However, by elementary manipulations, (28) becomes

$$|\Delta\alpha| = |\cos^{-1}(z \cos \alpha) - \alpha|,$$

where, temporarily,

$$z \equiv \cos \delta + \frac{\Delta\mu_3}{\Gamma}. \quad (30)$$

By considering the function $z \mapsto |\cos^{-1}(z \cos \alpha) - \alpha|$ for small α one verifies that $|\cos^{-1}(z \cos \alpha) - \alpha| \ll \alpha$ is equivalent to

$$0 < \frac{z-1}{\frac{1}{\cos \alpha} - 1} \ll 1 \quad \text{or} \quad 0 < \frac{1-z}{\frac{\cos 2\alpha}{\cos \alpha} - 1} \ll 1.$$

In any case this amounts to $|z-1| < O(\alpha^2)$ and since $\delta \ll \alpha$ one gets by (30) that $\Delta\mu_3 \ll O(\alpha^2)$. The point is that for momentum perturbations $\Delta\mu_1$, $\Delta\mu_3$ to be considered small the first must fall as α while the second must fall as α^2 . Since $|\Delta\mu_1| \gg |\Delta\mu_3|$ causes preq motion that is rotation about an axis perpendicular




Figure 2: Left: A preq incident from the lower left approaches its anti-preq incident from the lower right; a repulsive force is evident. Right: the same preq and anti-preq collide head on and break up into dipole pairs. The bars at the lower left indicate the diameter of the preqs.

to the direction of the preq, the effect is that the preqs can have momentum $O(\alpha)$ only for motions $O(\alpha)$ -close to great circle paths. To obtain preq motion along smaller circular paths on the sphere one must take $\Delta\mu_1 \approx \Delta\mu_3 = O(\alpha^2)$, which implies a much smaller $O(\alpha^2)$ momentum.

Since the maximum reasonable momentum perturbations are $O(\alpha)$ and the 4-vortex mass is $O(\alpha^2)$, the maximum 4-vortex angular velocity on the sphere is $O(1/\alpha)$. Thus greater velocities are available to smaller opening angles; the velocity available to a 4-vortex preq compared with its diameter is $O(1/\alpha^2)$.

A most interesting aspect of preqs with small opening angles is that more than one of them may be positioned on the sphere, all initially far apart (compared with their radii) from one another. Since the vortex-vortex interaction falls as the the vortex-vortex distance increases, as long as the separate preqs remain separated they will not strongly interact with one another, and they will maintain their separate identities. However, as they separately move on the sphere they may closely approach one another. When two or more preqs closely approach the above perturbation theory becomes inapplicable and they undergo an interaction, and may move apart from one another largely unchanged, or may be partially or completely destroyed. There is no theory of these interactions at this time but the main features, obtained by numerical simulation and illustrated in Figures (2) and (3), are as follows:

- Two preqs interact when their distance falls to lengths comparable with their diameters.
- Two preqs with vortices having the same sign are attracted to one another while if the vortices have opposite signs they repel one another. Thus identical preqs attract while a preq and its anti-preq (obtained from the first by changing the signs of each vortex) repel.
- The collision of two identical preqs, partially since those preqs attract, tends to result in two groups of four vortices representing a state of the vortex system far away from the relative equilibria p_e . In this sense the 4-vortex preqs are usually destroyed in same-sense vortex preq collisions.




Figure 3: Left: A preq incident from the lower left approaches an identical preq initially at rest in the center; and attractive force is evident. Right: a preq moving rightward and slightly upward collides with an identical preq moving leftward and slightly downward. The denser tracks at the center are the incident preqs. The preq incident from the left remains and rebounds back the way it came. The preq incident from the right has been nearly destroyed and is a four vortex motion far from the original relative equilibrium. Clearly visible in the interaction is the exchange of an outer vortex.

In such an interaction the two 4-vortex preqs usually exchange one of their outer vortices.

- A very energetic collision of a 4-vortex preq with its anti-preq usually results in the destruction of both preqs into vortex dipole pairs.
- The detailed results of an energetic or non-elastic 4-vortex preq collisions are extremely sensitive to initial condition, while the gross aspects of the collision (e.g. whether or not the collision is elastic) are relatively more robust.

Many a happy hour may be spent watching the antics of 4-vortex preqs colliding of the sphere and the nature of the 4-vortex interaction is the subject of ongoing investigations.

4 Transcription to the Planar System

As already noted (see Equations (8)), when restricted to sufficiently small regions of the sphere, the system of point vortices on the sphere reverts to the system of point vortices on the plane. This is pertinent for the study of small-opening-angle preq interactions, since the interactions occur only as the preq closely approach. Moreover, the planar case, being an example with noncompact symmetry group, is currently interesting, as the general focus of the Hamiltonian stability literature has been on the compact case, while the noncompact situation is known to have distinctive features, as shown by Leonard and Marsden [1997]. Consequently, I close this article with a short transcription of the above the the planar case; the results should be viewed as the $\alpha = 0$ limit of the spherical system.

The transcription is easy: through the pull back (7) of the sphere to the plane, the relative equilibria p_e with opening angle α have radii $\tilde{\alpha} = R \sin \alpha$,

and by (20) they rotate at angular velocity

$$\dot{\theta}_\alpha = \frac{\Gamma}{4\pi R^2 \sin^2 \alpha} \left(\frac{1}{N-1} + \cos \alpha \right) = \frac{N}{N-1} \frac{\Gamma}{4\pi \tilde{\alpha}^2} + O(\tilde{\alpha}^4), \quad (31)$$

while, from (13),

$$\Gamma_1 = \frac{-\Gamma}{(N-1) \cos \alpha} = \frac{-\Gamma}{N-1} + O(\tilde{\alpha}^2).$$

Deleting the higher order terms and renaming $\tilde{\alpha}$ to α , one expects planar relative equilibria with radii α , central vortex strength Γ , outer vortex strength $\Gamma/(N-1)$, and $\mathfrak{se}(2)$ generator $(\dot{\theta}_\alpha, 0)$, where

$$\dot{\theta}_\alpha = \frac{N}{N-1} \frac{\Gamma}{4\pi \alpha^2},$$

and it is easily verified that these are indeed relative equilibria of the planar system. Moreover, these relative equilibria occur for the vortex strength parameters Γ_n such that $\sum \Gamma_n = 0$, parameters for which, by (10), the momentum map J^{pl} is equivariant.

For my purpose, equivariance of the momentum map is important since, in the general, equivariance guarantees, for a relative equilibrium p_e with momentum μ_e and generator ξ_e , the *momentum-commutation relation* $\text{coad}_{\xi_e} \mu_e = 0$, and this is crucial for the perturbation theory of Patrick [1995]. On the other hand, in absence of equivariance one has the more complicated commutation relation

$$\text{coad}_{\xi_e} \mu_e = -\Sigma(\xi_e, \cdot) = -i_{\xi_e} \Sigma,$$

and the perturbation theory would have to be extended in a fundamental way to cover the nonequivariant case. That not being necessary, one calculates the nilpotent parts of the linearizations of the planar system at the relative equilibria above, to obtain, for the 3-vortex relative equilibria

$$N_\alpha = \frac{1}{2\pi \alpha^4} \begin{pmatrix} -\frac{3}{4} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

while for the 4-vortex relative equilibria the nilpotent part turns out to be

$$N_\alpha = \frac{1}{2\pi \alpha^2} \begin{pmatrix} -\frac{4}{3} & 0 & 0 \\ 0 & \frac{3}{4} & 0 \\ 0 & 0 & \frac{3}{4} \end{pmatrix},$$

Going to the Lagrangian viewpoint, and comparing with the Lagrangian of a 2-dimensional rigid body moving in the plane, one sees that the 3-vortex preq have infinite mass while the 4-vortex preq have inertia coefficient $-\frac{3}{2\pi} \alpha^2$ and mass $\frac{8\pi}{3} \alpha^2$.

References

- Abraham, R. and J. E. Marsden [1978]. *Foundations of Mechanics* (second ed.). Addison-Wesley.
- Adams, M. and T. S. Ratiu [1988]. The three point vortex problem: commutative and non-commutative integrability. In K. R. Meyer and D. Saari (Eds.), *Hamiltonian dynamical systems*, Volume 81 of *Contemporary Mathematics*, pp. 245–257. AMS.
- Channell, P. J. and F. P. Neri [1993]. An introduction to symplectic integrators. In J. E. Marsden, G. W. Patrick, and W. F. Shadwick (Eds.), *Integration algorithms and classical mechanics*, Volume 10 of *Fields Inst. Comm.*, pp. 45–58.
- Kidambi, R. and P. K. Newton [1998]. Motion of three point vortices on a sphere. *Physica D* 116, 143–175.
- Leonard, N. E. and J. E. Marsden [1997]. Stability and drift of underwater vehicle dynamics: mechanical systems with rigid motion symmetry. *Physica D* 105, 130–162.
- Lewis, D. and T. S. Ratiu [1996]. Rotating n -gon/ kn -gon vortex configurations. *J. Nonlin. Sc.* 6, 385–414.
- Marsden, J. E. and T. S. Ratiu [1994]. *Introduction to Mechanics and Symmetry*, Volume 17 of *Texts in Applied Mathematics*. Springer-Verlag.
- Patrick, G. W. [1992]. Relative equilibria in Hamiltonian systems: The dynamic interpretation of nonlinear stability on the reduced phase space. *J. Geom. Phys.* 9, 111–119.
- Patrick, G. W. [1995]. Relative equilibria of Hamiltonian systems with symmetry: linearization, smoothness, and drift. *J. Nonlin. Sc.* 5, 373–418.
- Patrick, G. W. [1998]. Dynamics near relative equilibria: Nongeneric momenta at a 1:1 group-reduced resonance. Submitted *Math. Z.*
- Pekarsky, S. and J. E. Marsden [1998]. Point vortices on the sphere: stability of relative equilibria. To appear *J. Math. Phys.*